

Aerobic Fitness Is Associated with Cerebral μ -Opioid Receptor Activation in Healthy Humans

AQ1 TIINA SAANIJOKI^{1,2}, TATU KANTONEN^{1,3}, LAURA PEKKARINEN^{1,4}, KARI KALLIOKOSKI¹, JUSSI HIRVONEN^{1,5}, TUULIA MALÉN¹, LAURI TUOMINEN^{1,6}, JETRO J. TUULARI^{1,7}, EVELIINA ARPONEN¹, PIRJO NUUTILA^{1,4}, and LAURI NUMMENMAA^{1,8}

¹Turku PET Centre, University of Turku, Turku, FINLAND; ²Turku BioImaging, University of Turku and Åbo Akademi University, Turku, FINLAND; ³Clinical Neurosciences, University of Turku and Turku University Hospital, Turku, FINLAND; ⁴Department of Endocrinology, Turku University Hospital, Turku, FINLAND; ⁵Department of Radiology, Turku University Hospital, Turku, FINLAND; ⁶The Royal's Institute of Mental Health Research, University of Ottawa, Ottawa, Ontario, CANADA; ⁷FinnBrain Birth Cohort Study, Turku Brain and Mind Center, Department of Clinical Medicine, University of Turku, Turku, FINLAND; and ⁸Department of Psychology, University of Turku, Turku, FINLAND

ABSTRACT

SAANIJOKI, T., T. KANTONEN, L. PEKKARINEN, K. KALLIOKOSKI, J. HIRVONEN, T. MALÉN, L. TUOMINEN, J. J. TUULARI, E. ARPONEN, P. NUUTILA, and L. NUMMENMAA. Aerobic Fitness Is Associated with Cerebral μ -Opioid Receptor Activation in Healthy Humans. *Med. Sci. Sports Exerc.*, Vol. 54, No. 7, pp. 00–00, 2022. **Introduction:** Central μ -opioid receptors (MORs) modulate affective responses to physical exercise. Individuals with higher aerobic fitness report greater exercise-induced mood improvements than those with lower fitness, but the link between cardiorespiratory fitness and the MOR system remains unresolved. Here we tested whether maximal oxygen uptake ($\dot{V}O_{2\text{peak}}$) and physical activity level are associated with cerebral MOR availability and whether these phenotypes predict endogenous opioid release after a session of exercise. **Methods:** We studied 64 healthy lean men who performed a maximal incremental cycling test for $\dot{V}O_{2\text{peak}}$ determination, completed a questionnaire assessing moderate-to-vigorous physical activity (MVPA; in minutes per week), and underwent positron emission tomography with [¹¹C]carfentanil, a specific radioligand for MOR. A subset of 24 subjects underwent additional positron emission tomography scan also after a 1-h session of moderate-intensity exercise and 12 of them also after a bout of high-intensity interval training. **Results:** Higher self-reported MVPA level predicted greater opioid release after high-intensity interval training, and both $\dot{V}O_{2\text{peak}}$ and MVPA level were associated with a larger decrease in cerebral MOR binding after aerobic exercise in the ventral striatum, orbitofrontal cortex, and insula. That is, more trained individuals showed greater opioid release acutely after exercise in brain regions especially relevant for reward and cognitive processing. Fitness was not associated with MOR availability. **Conclusions:** We conclude that regular exercise training and higher aerobic fitness may induce neuroadaptation within the MOR system, which might contribute to improved emotional and behavioral responses associated with long-term exercise. **Key Words:** OPIOID SYSTEM, POSITRON EMISSION TOMOGRAPHY (PET), BRAIN IMAGING, FITNESS, PHYSICAL ACTIVITY LEVEL

Habitual physical activity and cardiorespiratory fitness (CRF) are well-established modifiable lifestyle factors that promote brain health throughout the life span. Higher fitness and greater amounts of physical activity are linked with better cognitive functioning (1), lower levels of anxiety and depression (2), and reduced risk for neurodegenerative disease (3). These biological and psychological

benefits of exercise are paralleled in brain structure and function. Better fitness and higher physical activity levels are associated with higher gray (4,5) and white matter volume (4,6). In addition, several intervention studies have demonstrated that improved fitness positively affects brain volumes in older adults, especially in frontotemporal regions that are important for cognition and memory functions, and most susceptible to age-related brain atrophy (7,8). Moreover, higher aerobic fitness promotes efficient functional connectivity of multiple brain networks supporting cognitive control and memory functions (9,10).

Physical exercise also acutely affects the functioning of the brain's neuromodulatory systems, particularly the endogenous opioid system (11). Endogenous opioid system and especially μ -opioid receptors (MORs) are closely involved in processing reward, motivation, and emotions (12). They also have a central role in several physiological functions, such as pain processing (13) and stress regulation (14), and recent evidence links opioid system dysregulation with depressive and anxious symptoms (15). Therefore, the opioid system could potentially

AQ2 Address for correspondence: Tiina Saanijoki, Turku PET Centre c/o Turku University Hospital, Kiinamyllynkatu 4-6, 20520 Turku, Finland; E-mail: tiina.saanijoki@utu.fi.

Submitted for publication June 2021.

Accepted for publication February 2022.

Supplemental digital content is available for this article. Direct URL citations appear in the printed text and are provided in the HTML and PDF versions of this article on the journal's Web site (www.acsm-msse.org).

0195-9131/22/5407-0000/0

MEDICINE & SCIENCE IN SPORTS & EXERCISE®

Copyright © 2022 by the American College of Sports Medicine

DOI: 10.1249/MSS.0000000000002895

mediate the psychological benefits of regular exercise, such as anxiety reduction and improved mood.

Animal studies investigating the effects of regular exercise training on the opioid system have found elevated β -endorphin and met-enkephalin levels in the periaqueductal gray area and rostral ventromedial medulla after 5 wk of treadmill running (16) and shown that chronic exercise, in comparison with short-term exercise or no exercise, decreases MOR expression (17) and overall MOR availability in rat brain (18). Exercising rats also show reduced sensitivity to antinociceptive effects of exogenous opioid agonists such as morphine, which may indicate downregulation of MORs resulting from increased endogenous opioid concentrations elevated by regular exercise training (19,20). Human neuroimaging studies have demonstrated that a single bout of moderate-to-vigorous exercise stimulates endogenous opioid release in the brain, which is associated with affective responses induced by exercise (21–23). Taken together, converging evidence from animal and human studies suggests that regular exercise training might induce neuroadaptation within the central MOR system, subsequently contributing to improvements in mood and stress regulation. However, *in vivo* evidence from humans is currently lacking.

Here we investigated whether individual differences in baseline MOR availability are associated with CRF and habitual physical activity levels in healthy young men. We used *in vivo* positron emission tomography (PET) imaging with the highly selective MOR agonist ligand [^{11}C]carfentanil. We coupled MOR data with measurement of peak oxygen consumption ($\dot{V}\text{O}_{2\text{peak}}$), an objective and direct measure of CRF, and with self-reported physical activity questionnaires. To test whether higher fitness and physical activity levels influence the capacity of acute exercise to activate the MOR system, we also studied a subset of participants with [^{11}C]carfentanil PET after a session of high-intensity interval training (HIIT) and after a 1-h session of aerobic exercise. Based on previous human and animal research, we hypothesized that higher fitness and physical activity levels would be negatively associated with cerebral MOR availability in the brain's reward circuits and positively associated with gray matter (GM) volume. In addition, we predicted that $\dot{V}\text{O}_{2\text{peak}}$ and self-reported physical activity would be associated with exercise-induced changes in MOR availability.

METHODS

Subjects. The Ethics Committee of the Hospital District of Southwest Finland approved the study protocol, and the study was conducted in accordance with the Declaration of Helsinki. The sample size was determined by *a priori* power analysis based on our prior neuroreceptor PET studies (24,25), which suggested that, with expected effect size of $r = 0.45$, a sample size of 45 would be sufficient for establishing the predicted effects at power of 0.95. Sixty-four male adults with a variable exercise background were enrolled in the study (Table 1). All subjects signed ethics committee–approved informed consent forms. They were recruited via Internet discussion forums, traditional bulletin boards, university-hosted email

TABLE 1. Subject characteristics ($n = 64$).

	Mean (SD)	Range
Age (yr)	25.4 (4.6)	20–36
BMI ($\text{kg}\cdot\text{m}^{-2}$)	24.1 (2.8)	18.5–31.0
Total physical activity ($\text{min}\cdot\text{wk}^{-1}$)	291 (165)	0–870
MVPA ($\text{min}\cdot\text{wk}^{-1}$)	160 (113)	0–870
$\dot{V}\text{O}_{2\text{max}}$ ($\text{mL}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$)	44.5 (7.9)	25.8–61.7

lists, and newspaper advertisements. The exclusion criteria were poor compliance, a history of or current neurological or psychiatric disease, use of tobacco products or medication affecting the central nervous system, current or past excessive alcohol or substance abuse, and standard PET and magnetic resonance imaging (MRI) exclusion criteria. Laboratory tests, urinalysis, and an ECG were obtained to assess health and the absence of psychoactive drugs. These data have originally been collected in clinical trials EXEBRAIN (NCT02615756; $n = 24$) and PROSPECT (NCT03106688; $n = 40$).

Physical activity and aerobic fitness measurements. Self-reported physical activity was assessed with a questionnaire where participants rated the frequency (days per week) and duration (hours and minutes per week) of moderate-to-vigorous physical activity (MVPA) and other physical activity during the last 3 months. CRF was assessed as $\dot{V}\text{O}_{2\text{peak}}$, which was determined in a maximal exercise test performed on a cycle ergometer starting at 40–50 W and followed by an increase of 30 W in every 2 min until volitional exhaustion. Ventilation and gas exchange were measured (Jaeger Oxycon Pro; VIASYS Healthcare) and reported as the mean value per minute. The highest 1-min mean value of oxygen consumption was expressed as the $\dot{V}\text{O}_{2\text{peak}}$.

PET data acquisition. We measured MOR availability with the agonist radioligand [^{11}C]carfentanil that has a high affinity for MORs. Radioligand syntheses for the EXEBRAIN and PROSPECT trials have been described previously (22,26). Subjects refrained from exercise for at least 24 h and fasted for at least 2 h before scanning. Data were acquired with the 3 T Philips Ingenuity TF PET/MR (Philips Healthcare, Cleveland, OH) scanner or PET/CT (GE Discovery VCT PET/CT, GE Healthcare (General Electric Medical Systems, Milwaukee, WI)) at Turku PET Centre. Data acquisition started concomitantly with the intravenous radioligand bolus injection ($M = 250 \text{ MBq}$, $SD = 13 \text{ MBq}$), and cerebral radioactivity was measured for 51 min. Data were corrected for dead time, decay, and measured photon attenuation.

PET challenge paradigm for exercise-induced opioid release. A subset of participants ($n = 24$; Table S1, Supplemental Digital Content, <http://links.lww.com/MSS/C538>) underwent an additional PET scan after a 1-h session of moderate-intensity cycling exercise on a separate day, and 12 of these participants underwent a PET scan also after a session of HIIT; the exercise protocols and opioid release data have been reported previously (22). The order of the exercise/rest PET studies was randomized and counterbalanced for these participants. Emotional reactions to physical exercise were measured with the Positive Affect and Negative Affect Schedule (27).

MRI acquisition. Anatomical MR images were acquired for VBM and for preprocessing the PET images with the 3 T **AQ3**

T1




FIGURE 1— $\dot{V}O_{2\text{peak}}$ correlated negatively with age (A) and BMI (B). Age correlated positively with BMI (C). $\dot{V}O_{2\text{peak}}$ was positively associated with change in positive affect after aerobic exercise (D).

Philips Ingenuity TF PET/MR scanner using a T1-weighted sequence with 1-mm³ resolution (repetition time, 8.1 ms; echo time, 3.7 ms; flip angle, 7°; scan time, 263 s). Complementary voxel-based morphometric analyses on the association between aerobic fitness, physical activity, and cerebral density are described in Supplemental Digital Content, <http://links.lww.com/MSS/C538>.

PET data preprocessing and analysis. PET data were processed with the automated Magia pipeline (28) (<https://github.com/tkjarjal/magia>). Processing began with motion correction of the PET data followed by coregistration of the PET and MR images. Magia uses FreeSurfer (<http://surfer.nmr.mgh.harvard.edu/>) to define the regions of interest (ROIs) as well as the reference regions. The ROI-wise kinetic modeling was based on the extraction of ROI-wise time–activity curves. The PET images were slightly smoothed using Gaussian kernel (2-mm full width at half maximum) to increase the signal-to-noise ratio before model fitting. Parametric images were spatially normalized to MNI space and finally smoothed using a Gaussian kernel (full width at half maximum = 6 mm). [¹¹C]carfentanil binding was quantified by binding potential (BP_{ND}), which is the ratio of specific binding to nondisplaceable binding in the tissue (29). The occipital cortex was used as the reference region (30). Because of technical problems with the PET scanner, the PET data after one aerobic exercise scan and one HIIT scan were subsequently found invalid and were excluded from the analysis.

Statistical analysis. The effects of $\dot{V}O_{2\text{peak}}$ and self-reported physical activity on i) MOR availability, ii) MOR activity after physical exercise, and iii) GM densities were assessed in SPM12 (<http://www.fil.ion.ucl.ac.uk/spm/>) using a linear regression model with body mass index (BMI) and PET scanner (for baseline PET data) as a covariate. The statistical threshold was set at $P < 0.05$, false discovery rate (FDR)

corrected at cluster level. Atlas-based ROIs were generated in the MOR-rich regions in the brain (31) (amygdala, hippocampus, ventral striatum, dorsal caudate, thalamus, insula, orbitofrontal cortex, anterior cingulate cortex, middle cingulate cortex, and posterior cingulate cortex) using AAL (32) and Anatomy (33) toolboxes. Mean regional [¹¹C]carfentanil BP_{ND} was extracted for each region, and the averaged ROI data were analyzed with R statistical software (<https://cran.r-project.org>) (34). Complementary hierarchical Bayesian analyses are described in Supplemental Digital Content, <http://links.lww.com/MSS/C538>.

RESULTS

$\dot{V}O_{2\text{peak}}$ is associated with physical activity level, age, and BMI. $\dot{V}O_{2\text{peak}}$ was positively associated with self-reported physical activity ($r = 0.45$, $P < 0.01$) and negatively associated with age ($r = -0.37$, $P < 0.01$; Fig. 1A) and BMI (F1) ($r = -0.46$, $P < 0.01$; Fig. 1B). Age was positively associated with BMI ($r = 0.38$, $P < 0.01$; Fig. 1C).

Higher MVPA level predicts a larger decrease in MOR availability after HIIT. We recently showed that HIIT significantly decreased MOR binding in human brain, indicative of endogenous opioid release (22). Here we tested whether exercise-induced changes in BP_{ND} would be associated with self-reported physical activity or $\dot{V}O_{2\text{peak}}$, indicative of exercise habit-dependent MOR activation. We found a negative association between MVPA level and changes in BP_{ND} after HIIT (Fig. 2), such that higher MVPA level was associated with a larger decrease in BP_{ND} after HIIT. This effect was observed in the ventral and dorsal striatum, hippocampus, left amygdala, thalamus, cingulate cortex, insular cortex, somatosensory cortex, temporal areas, and orbitofrontal cortex. $\dot{V}O_{2\text{peak}}$ and total minutes of self-reported physical activity




FIGURE 2—Negative association between self-reported MVPA level and changes in BP_{ND} after HIIT session in a subset of 11 participants. The data are thresholded at $P < 0.05$, FDR corrected at the cluster level. Scatterplot shows the corresponding association (least squares regression line with 95% confidence interval) in the orbitofrontal cortex.




FIGURE 3—Higher fitness level predicted higher exercise-induced opioid release after moderate-intensity exercise, as indicated by a negative association between changes in BP_{ND} after 1-h session of aerobic exercise and $\dot{V}O_{2peak}$ (A) and self-reported MVPA level (B). The data are thresholded at $P < 0.05$, FDR corrected at the cluster level. Scatterplots show the corresponding association (least squares regression line with 95% confidence interval) in the putamen.

were not associated with HIIT-induced BP_{ND} changes in this small sample of 11 participants.

Higher $\dot{V}O_{2peak}$ and MVPA level predict a larger decrease in MOR availability after aerobic exercise.

MOR availability shows notable variation between individuals after aerobic exercise, such that [^{11}C]carfentanil BP_{ND} decreases in some individuals and increases in others (22). Here

[F3] we found that both $\dot{V}O_{2peak}$ (Fig. 3A) and self-reported MVPA level (Fig. 3B) were negatively associated with exercise-induced change in BP_{ND} , such that higher $\dot{V}O_{2peak}$ and higher MVPA level predicted a larger decrease in BP_{ND} after exercise. This effect was observed in the ventral and dorsal striatum, left hippocampus, left thalamus, insular cortex, somatosensory cortex, temporal areas, and orbitofrontal cortex. No associations were found between total self-reported physical activity and exercise-induced change in BP_{ND} .

We previously reported enhanced mood responses after aerobic exercise (22). Here, we found a positive association between $\dot{V}O_{2peak}$ and change in positive affect as measured with the Positive Affect and Negative Affect Schedule before and after aerobic

exercise ($r = 0.59$, $P < 0.01$; Fig. 1D), indicating that higher $\dot{V}O_{2peak}$ was associated with greater mood improvement.

Association of aerobic fitness and physical activity level on MOR availability.

We next tested whether baseline differences in aerobic fitness are associated with MOR availability. Full-volume analysis showed a negative association between $\dot{V}O_{2peak}$ and baseline MOR availability (BP_{ND}) in a large cluster extending to both hemispheres from the frontal lobe to the parieto-occipital sulcus (Fig. 4). Significant associ-

[F4] ations were also observed in bilateral putamen, thalamus, insula, and temporal cortices. In addition, the ROI analysis revealed significant associations in orbitofrontal and middle cingulate cortices ($P < 0.05$). Comparable analysis where MOR availability was predicted with self-reported physical activity yielded similar effects, but only when BMI was not controlled for in the model (data not shown). Because gross brain atrophy is negligible in healthy subjects younger than 37 yr, partial volume effects due to atrophy are unlikely to significantly bias our results. Therefore, we did not correct the data for partial volume effects.




FIGURE 4—Negative association between $\dot{V}O_{2peak}$ and baseline [^{11}C]carfentanil BP_{ND} . The data are thresholded at $P < 0.05$, FDR corrected at the cluster level. Scatterplots show the corresponding association (least squares regression line with 95% confidence interval) in representative anatomical ROIs.

Aging has regionally specific effects on MOR availability (31). Although our subjects had a relatively narrow age range ($M = 25.4$ yr, $SD = 4.6$ yr), we nevertheless wanted to statistically control for potential aging-dependent effects in the MOR availability. When age was entered in the analysis as a covariate, no associations were observed between MOR availability and $\dot{V}O_{2\text{peak}}$ or self-reported physical activity at the *a priori* statistical threshold. The unthresholded result files are available in Neurovault (<https://neurovault.org/collections/PTWTWOGG/>). $\dot{V}O_{2\text{peak}}$ showed a positive association with GM density when controlling for age and BMI (Fig. S1, Supplemental Digital Content, <http://links.lww.com/MSS/C538>). These findings are described in detail in Supplemental Digital Content, <http://links.lww.com/MSS/C538>.

DISCUSSION

The present findings indicate that aerobic fitness and habitual physical activity level have a previously unrecognized role in brain opioid signaling. Higher self-reported MVPA level was associated with a larger decrease in cerebral MOR binding after HIIT, and both higher MVPA level and higher CRF, as measured by $\dot{V}O_{2\text{peak}}$, were associated with a larger decrease in cerebral MOR binding after moderate-intensity aerobic exercise. These effects were observed in widespread cortical and subcortical areas, most notably in the anterior cingulate cortex, insula, orbitofrontal cortex, and ventral striatum. In other words, more trained individuals showed greater acute opioid release after exercise in brain regions involved in reward and cognitive processing. Fitness and physical activity level were not associated with baseline BP_{ND} . Conversely, $\dot{V}O_{2\text{peak}}$ was positively associated with GM density in several brain regions, including the medial and lateral frontal and orbitofrontal cortices, cingulate cortex, and striatum. Taken together, these data suggest that aerobic fitness and exercise training at moderate-to-vigorous intensity may modulate cerebral MOR function, which might be an important pathway regulating exercise habits, and that aerobic fitness and physically active lifestyle are associated with neurobiological markers of brain health (i.e., GM density) not only in older adults but also in early adulthood.

Higher MVPA level and aerobic fitness predict a greater decrease in MOR binding after exercise. Previous research has shown that exercise intensity modulates opioid action in the brain (22,23). High-intensity exercise induces a robust opioid release, whereas moderate-intensity exercise results in decreased MOR availability in some and increased in other individuals (22). Here we report, for the first time, that higher MVPA level is associated with higher cerebral opioid release after a bout of HIIT and that both higher $\dot{V}O_{2\text{peak}}$ and MVPA levels are associated with larger decrease in cerebral MOR binding after moderate-intensity aerobic exercise. Together, these findings suggest that aerobic fitness and physical activity level may shape opioidergic response after exercise. The dependency on prior aerobic fitness may also explain why moderate-intensity exercise does not result in net opioid release when individuals with different fitness statuses

are assessed together (22). The present findings go beyond past reports, which have only examined the association of training status on peripheral opioid concentrations after exercise. Higher circulating β -endorphin levels have been found in well-trained athletes, in comparison with untrained individuals, after a graded exercise test (35) and after a bout of supramaximal exercise (36). In contrast, high-intensity cycling (70% and 80% of maximal oxygen uptake ($\dot{V}O_{2\text{max}}$)) resulted in a similar increase in plasma β -endorphin concentration in both trained and untrained individuals, whereas moderate-intensity cycling (60% of $\dot{V}O_{2\text{max}}$) showed no effect on plasma β -endorphin concentration (37). Although it has been suggested that training-induced adaptation within the opioid system could increase the response capacity to extreme exercise stress (36), peripheral opioid levels probably do not mirror those of the brain (38) and thus limits reasonable comparison of these studies.

We observed greater decreases in MOR binding in more trained individuals after exercise in the ventral and dorsal striatum, left amygdala, hippocampus, thalamus, insular cortex, somatosensory cortex, temporal areas, and orbitofrontal cortex, and across the cingulate. MORs in these regions are closely involved in processing both nociceptive and hedonic signals (12) as well as modulating decision making and cognitive control (39). Spatially more widespread correlation was observed between training variables and the change in MOR binding after HIIT than after aerobic exercise, although unequal sample sizes may confound their comparison. Although both increased and decreased MOR binding were observed after moderate-intensity aerobic exercise, it should be noted that the within-subject variability in MOR responses after aerobic exercise far exceeds test-retest reproducibility. For example, in the thalamus and anterior cingulate cortex, the test-retest variabilities were 350% and 60% higher, respectively, in the present study in comparison to the prior study examining [^{11}C]carfentanil test-retest reliability (40).

After HIIT, higher training level was associated with greater opioid release in the thalamus, insula, left amygdala, and anterior cingulate cortex—brain regions known to process sensory and affective dimensions of pain (41–43). Enhanced MOR activation in these regions may improve exercise tolerance in more trained individuals by reducing pain and discomfort levels and thus help to sustain the metabolic and mental demands of high-intensity exercise. Higher fitness level was recently also found to be associated with larger pain tolerance after high-intensity but not after low-intensity exercise (44), indicating a relationship between fitness, exercise intensity, and opioid modulation. Moreover, the insular cortex is the main brain site responsible for the awareness of subjective feelings from the body (45), and it has been shown to play a central role especially in fatiguing exercise. The magnitude of insular activation varies with the intensity of exercise and associates with subjective ratings of perceived exertion (46). Interestingly, opioid modulation has shown to influence perceived exertion and exercise capacity (47). In addition to insula, prefrontal areas and anterior cingulate cortex have been

proposed to be implicated in the capacity to tolerate high levels of physical exertion and the determination of exercise termination (48). Consequently, greater HIIT-induced opioid release in more trained individuals may modify subjective perception of pain and fatigue and may thereby contribute to greater tolerance of higher exercise intensities and improved exercise performance.

The improved capacity of more trained individuals to activate the MOR system in response to exercise may also represent a neurobiological adaptation process induced by regularly repeated exercise bouts. Thus, improved MOR activation capacity could reinforce the adoption and maintenance of new exercise routines and, speculatively, underlie the rapid positive affective adaptation observed already within a few training sessions of both moderate- and high-intensity exercise in previously sedentary individuals (49,50). In line with this, we found that participants with higher CRF experienced greater improvements in mood after the aerobic exercise. This accords with previous studies reporting that better fitness level (51) and regular exercise participation are associated with more positive affective responses (52,53) and enhanced anxiety relief (52,54) after a bout of exercise. We propose that greater opioid release could explain enhanced emotional and antinociceptive responses reported by people with higher exercise levels and, thus, bear implications in exercise adoption and engagement (55), yet this idea remains to be determined in future studies.

Association of aerobic fitness and physical activity level with MOR availability. We found that $\dot{V}O_{2\text{peak}}$ and self-reported physical activity were negatively correlated with baseline [^{11}C]carfentanil BP_{ND} , suggesting that higher levels of regular exercise promoting better aerobic fitness may induce neuroadaptation within the endogenous opioid system. However, these associations were no longer statistically significant when age was introduced as a covariate in the model between MOR availability and aerobic fitness or self-reported physical activity, despite our subjects' narrow age range (20–36 yr). Indeed, recent work has shown a prominent effect of age on MORs, especially before the age of 40 yr (31). Aging increases MOR availability in frontotemporal areas and decreases it in the thalamus and nucleus accumbens (31). Thus, pure age effects would unlikely explain our findings, for example, in the thalamus. Given that also CRF is markedly influenced by age, physical activity level, and BMI (56,57), it remains inconclusive whether the presently observed effects on MOR availability reflect mere age-specific effects or joint effects between fitness and age. Age-dependent MOR downregulation may, in fact, be propelled by declining physical fitness, yet this speculation warrants further research in more controlled subject cohorts allowing for differentiation between the intertwined effects of age, CRF, physical activity level, and BMI on MOR availability.

Prior animal studies have established a relationship between habitual physical activity and MOR expression (17,18). Chronic exercise, in comparison with short-term exercise or no exercise, decreases MOR expression (17) and overall MOR binding in rat brain (18), demonstrating a causal link between exercise and MOR binding. Reduced MOR binding

may reflect downregulation of MORs, increased opioid tone, or, after competition between endogenous opioids and the radioligand, a combination of the two. Animal studies suggest that regular exercise training increases tonic endogenous opioid levels. In rats, exercise training for 5 to 8 wk increases basal β -endorphin concentration in cerebrospinal fluid (58) and plasma (59) and elevates both β -endorphin and met-enkephalin levels in the periaqueductal gray area and rostral ventromedial medulla (16). Such rise in tonic opioid levels is also associated with altered pain processing, as regular exercise training increases nociceptive threshold in rats (60) and reverses measures of pain in animal models of chronic pain (16,61,62), and as these effects can be reversed by an opioid receptor antagonist naloxone (16,60–62). Exercise-induced opioidergic pain modulation has also been demonstrated in humans. Six-week aerobic exercise intervention decreased pain intensity and interference in low active patients with chronic low back pain, and participants with the highest number of minutes in the target exercise intensity zone exhibited the greatest increases in endogenous opioid analgesia as indexed by naloxone–placebo condition differences in evoked pain responses (63). This suggests a dose–response effect so that higher intensity and amount of exercise may generate greater adaptation of the MOR system. Chronic exercise has also been demonstrated to result in decreased sensitivity to analgesic effects of exogenous opioid agonists such as morphine both in humans (64) and animals (19,20), which indicates that regular exercise may induce cross-tolerance to exogenously administered opioid agonists because of greater concentrations of endogenous opioid peptides. Altogether, these studies indicate a role of habitual physical activity in the MOR system, which may contribute to long-term adaptations of various physiological and behavioral responses associated with regular exercise, such as improved mood and pain and stress regulation.

Exercise-induced affective and antinociceptive responses may also be modulated by other neurotransmitter systems, such as the endocannabinoid system (65). Plasma endocannabinoid levels increase peripherally after acute aerobic exercise in humans (65–68) independent of physical activity level (67), whereas long-term aerobic exercise decreases circulating endocannabinoid levels (69,70). Given the close interaction of opioid and endocannabinoid systems in the reward and emotion processing brain pathways (71–73), further studies should also examine central endocannabinoid signaling in exercise settings.

Limitations. The interpretation of [^{11}C]carfentanil PET studies is challenging. High-intensity exercise led to a significant decrease in MOR binding, whereas moderate-intensity exercise resulted in both decreased and increased MOR binding. According to the competition principle, reduced [^{11}C]carfentanil BP_{ND} is typically interpreted as evidence of increased endogenous opioid release. Similarly, increased BP_{ND} has been suggested to mirror MOR “deactivation” as an acute decrease in synaptic endogenous opioids (42,74). However, because BP_{ND} is a composite measure that does not differentiate between receptor density and affinity, radioligand binding

can also be affected by receptor trafficking and changes in binding affinity. Although we cannot directly specify which interpretation is most appropriate, our findings nevertheless are consistent with the modulatory role of aerobic fitness and physical activity level in MOR responses after both high-intensity and moderate-intensity exercises. Next, our data were sampled from two distinct projects using two different PET scanners. Although the outcome measure BP_{ND} theoretically controls for minor differences in scanner signal-to-noise ratios (75), comparable PET scanning protocols may yield different BP_{ND} estimates across scanners. Our recent study comparing BP_{ND} values of various PET ligands across PET scanners, however, suggests that the BP_{ND} values obtained from PET/CT and PET/MRI are sufficiently comparable for [¹¹C]carfentanil (76). Even though we corrected for potential scanner-related biases in the analyses, a possible confounding effect of different scanners remains in the analyses. Furthermore, because females and males may have differential neurochemical responses to exercise (77) and differences in brain MOR availability (31), the findings may not be generalizable to females. Finally, because of underpowered sample, this first demonstration of the role of fitness and training measures on exercise-induced opioid release should be considered as preliminary evidence.

CONCLUSIONS

We conclude that higher training status is associated with greater reductions in MOR availability after a bout of

high-intensity exercise, suggesting greater exercise-induced opioid release in more trained individuals. Higher MVPA level and higher aerobic fitness also predict exercise-induced changes in MOR binding after a bout of aerobic exercise, indicating a role of habitual physical activity in MOR modulation also at moderate exercise intensities. Aerobic fitness was positively associated with GM density but not associated with baseline MOR availability. Altogether, our findings suggest that improving aerobic fitness by regular physical activity of moderate to high intensity may induce neuroadaptation within the MOR system by improving exercise-induced opioid functioning, which may further modulate physiological and behavioral responses governed by the opioid system.

This study was supported by the Academy of Finland (grants nos. 251125, 251399, 256470, 265917, 281440, 283319, 304385, and 332225) and Sigrid Juselius foundation. The authors thank Emil Aaltonen Foundation, Finnish Cultural Foundation (Southwest Finland Fund), and Jenny and Antti Wihuri Foundation for personal grants to T. K., and Päivikki and Sakari Sohlberg Foundation and the State research funding of the expert responsibility area of the Tyks Turku University Hospital for personal grants to T. M.

Earlier version of this manuscript has been posted on preprint server medRxiv: <https://medrxiv.org/cgi/content/short/2020.12.13.20247627v1>.

The authors declare no conflict of interest. The results of the present study do not constitute endorsement by the American College of Sports Medicine. The results of the study are presented clearly, honestly, and without fabrication, falsification, or inappropriate data manipulation.

REFERENCES

1. Zhu N, Jacobs DR, Schreiner PJ, et al. Cardiorespiratory fitness and cognitive function in middle age: the CARDIA study. *Neurology*. 2014;82(15):1339–46.
2. Baumeister SE, Leitzmann MF, Bahls M, et al. Associations of leisure-time and occupational physical activity and cardiorespiratory fitness with incident and recurrent major depressive disorder, depressive symptoms, and incident anxiety in a general population. *J Clin Psychiatry*. 2017;78(1):e41–7.
3. DeFina LF, Willis BL, Radford NB, et al. The association between midlife cardiorespiratory fitness levels and later-life dementia: a cohort study. *Ann Intern Med*. 2013;158(3):162–8.
4. Boots EA, Schultz SA, Oh JM, et al. Cardiorespiratory fitness is associated with brain structure, cognition, and mood in a middle-age cohort at risk for Alzheimer's disease. *Brain Imaging Behav*. 2015;9(3):639–49.
5. Raichlen DA, Klimentidis YC, Bharadwaj PK, Alexander GE. Differential associations of engagement in physical activity and estimated cardiorespiratory fitness with brain volume in middle-age to older adults. *Brain Imaging Behav*. 2020;14(5):1994–2003.
6. Johnson NF, Kim C, Clasey JL, Bailey A, Gold BT. Cardiorespiratory fitness is positively correlated with cerebral white matter integrity in healthy seniors. *Neuroimage*. 2012;59(2):1514–23.
7. Colcombe SJ, Erickson KI, Scalf PE, et al. Aerobic exercise training increases brain volume in aging humans. *J Gerontol A Biol Sci Med Sci*. 2006;61(11):1166–70.
8. Erickson KI, Voss MW, Prakash RS, et al. Exercise training increases size of hippocampus and improves memory. *Proc Natl Acad Sci U S A*. 2011;108(7):3017–22.
9. Voss MW, Weng TB, Burzynska AZ, et al. Fitness, but not physical activity, is related to functional integrity of brain networks associated with aging. *Neuroimage*. 2016;131:113–25.
10. Talukdar T, Nikolaidis A, Zwilling CE, et al. Aerobic fitness explains individual differences in the functional brain connectome of healthy young adults. *Cereb Cortex*. 2018;28(10):3600–9.
11. Greenwood BN, Foley TE, Le TV, et al. Long-term voluntary wheel running is rewarding and produces plasticity in the mesolimbic reward pathway. *Behav Brain Res*. 2011;217(2):354–62.
12. Nummenmaa L, Tuominen L. Opioid system and human emotions. *Br J Pharmacol*. 2018;175(14):2737–49.
13. Corder G, Castro DC, Bruchas MR, Scherrer G. Endogenous and exogenous opioids in pain. *Annu Rev Neurosci*. 2018;41(1):453–73.
14. Ribeiro SC, Kennedy SE, Smith YR, Stohler CS, Zubieta JK. Interface of physical and emotional stress regulation through the endogenous opioid system and mu-opioid receptors. *Prog Neuropsychopharmacol Biol Psychiatry*. 2005;29(8):1264–80.
15. Nummenmaa L, Karjalainen T, Isojärvi J, et al. Lowered endogenous mu-opioid receptor availability in subclinical depression and anxiety. *Neuropsychopharmacology*. 2020;45(11):1953–9.
16. Stagg NJ, Mata HP, Ibrahim MM, et al. Regular exercise reverses sensory hypersensitivity in a rat neuropathic pain model: role of endogenous opioids. *Anesthesiology*. 2011;114(4):940–8.
17. de Oliveira MSR, da Silva Fernandes MJ, Scorza FA, et al. Acute and chronic exercise modulates the expression of MOR opioid receptors in the hippocampal formation of rats. *Brain Res Bull*. 2010;83(5):278–83.
18. Arida RM, Gomes da Silva S, de Almeida AA, et al. Differential effects of exercise on brain opioid receptor binding and activation in rats. *J Neurochem*. 2015;132(2):206–17.
19. Smith MA, Lyle MA. Chronic exercise decreases sensitivity to mu opioids in female rats: correlation with exercise output. *Pharmacol Biochem Behav*. 2006;85(1):12–22.

20. Smith MA, Yancey DL. Sensitivity to the effects of opioids in rats with free access to exercise wheels: mu-opioid tolerance and physical dependence. *Psychopharmacology (Berl)*. 2003;168(4):426–34.
21. Boecker H, Sprenger T, Spilker ME, et al. The runner's high: opioidergic mechanisms in the human brain. *Cereb Cortex*. 2008;18(11):2523–31.
22. Saanijoki T, Tuominen L, Tuulari JJ, et al. Opioid release after high-intensity interval training in healthy human subjects. *Neuropsychopharmacology*. 2018;43(2):246–54.
23. Hiura M, Sakata M, Ishii K, et al. Central μ -opioidergic system activation evoked by heavy and severe-intensity cycling exercise in humans: a pilot study using positron emission tomography with ¹¹C-carfentanil. *Int J Sports Med*. 2017;38(1):19–26.
24. Karjalainen T, Tuominen L, Manninen S, et al. Behavioural activation system sensitivity is associated with cerebral μ -opioid receptor availability. *Soc Cogn Affect Neurosci*. 2016;11(8):1310–6.
25. Nummenmaa L, Manninen S, Tuominen L, et al. Adult attachment style is associated with cerebral μ -opioid receptor availability in humans. *Hum Brain Mapp*. 2015;36(9):3621–8.
26. Kantonen T, Pekkarinen L, Karjalainen T, et al. Obesity risk is associated with altered cerebral glucose metabolism and decreased μ -opioid and CB₁ receptor availability. *Int J Obes (Lond)*. 2022;46:400–7.
27. Watson D, Clark LA, Tellegen A. Development and validation of brief measures of positive and negative affect: the PANAS scales. *J Pers Soc Psychol*. 1988;54(6):1063–70.
28. Karjalainen T, Tuisku J, Santavirta S, et al. Magia: robust automated image processing and kinetic modeling toolbox for PET neuroinformatics. *Front Neuroinform*. 2020;14:3.
29. Innis RB, Cunningham VJ, Delforge J, et al. Consensus nomenclature for *in vivo* imaging of reversibly binding radioligands. *J Cereb Blood Flow Metab*. 2007;27(9):1533–9.
30. Frost JJ, Douglass KH, Mayberg HS, et al. Multicompartmental analysis of [¹¹C]-carfentanil binding to opiate receptors in humans measured by positron emission tomography. *J Cereb Blood Flow Metab*. 1989;9(3):398–409.
31. Kantonen T, Karjalainen T, Isojärvi J, et al. Interindividual variability and lateralization of μ -opioid receptors in the human brain. *Neuroimage*. 2020;217:116922.
32. Tzourio-Mazoyer N, Landeau B, Papathanassiou D, et al. Automated anatomical labeling of activations in SPM using a macroscopic anatomical parcellation of the MNI MRI single-subject brain. *Neuroimage*. 2002;15(1):273–89.
33. Eickhoff SB, Stephan KE, Mohlberg H, et al. A new SPM toolbox for combining probabilistic cytoarchitectonic maps and functional imaging data. *Neuroimage*. 2005;25(4):1325–35.
34. Rosseel Y. Lavaan: an R package for structural equation modeling. *J Stat Softw*. 2012;48(1):1–36.
35. de Diego Acosta AM, García JC, Fernández-Pastor VJ, Perán S, Ruiz M, Guirado F. Influence of fitness on the integrated neuroendocrine response to aerobic exercise until exhaustion. *J Physiol Biochem*. 2001;57(4):313–20.
36. Farrell PA, Kjaer M, Bach FW, Galbo H. Beta-endorphin and adrenocorticotropin response to supramaximal treadmill exercise in trained and untrained males. *Acta Physiol Scand*. 1987;130(4):619–25.
37. Goldfarb AH, Hatfield BD, Potts J, Armstrong D. Beta-endorphin time course response to intensity of exercise: effect of training status. *Int J Sports Med*. 1991;12(3):264–8.
38. Dietrich A, McDaniel WF. Endocannabinoids and exercise. *Br J Sports Med*. 2004;38(5):536–41.
39. van Steenbergen H, Eikemo M, Leknes S. The role of the opioid system in decision making and cognitive control: a review. *Cogn Affect Behav Neurosci*. 2019;19(3):435–58.
40. Hirvonen J, Aalto S, Hagelberg N, et al. Measurement of central μ -opioid receptor binding *in vivo* with PET and [¹¹C]carfentanil: a test–retest study in healthy subjects. *Eur J Nucl Med Mol Imaging*. 2009;36(2):275–86.
41. Bencherif B, Fuchs PN, Sheth R, Dannals RF, Campbell JN, Frost JJ. Pain activation of human supraspinal opioid pathways as demonstrated by [¹¹C]-carfentanil and positron emission tomography (PET). *Pain*. 2002;99(3):589–98.
42. Zubieta JK, Smith YR, Bueller JA, et al. Mu-opioid receptor-mediated antinociceptive responses differ in men and women. *J Neurosci*. 2002;22(12):5100–7.
43. Zubieta JK, Smith YR, Bueller JA, et al. Regional mu opioid receptor regulation of sensory and affective dimensions of pain. *Science*. 2001;293(5528):311–5.
44. Schmitt A, Wallat D, Stangier C, Martin JA, Schlesinger-Irsch U, Boecker H. Effects of fitness level and exercise intensity on pain and mood responses. *Eur J Pain*. 2020;24(3):568–79.
45. Craig AD. How do you feel? Interoception: the sense of the physiological condition of the body. *Nat Rev Neurosci*. 2002;3(8):655–66.
46. Williamson JW, McColl R, Mathews D, Ginsburg M, Mitchell JH. Activation of the insular cortex is affected by the intensity of exercise. *J Appl Physiol*. 1999;87(3):1213–9.
47. Sgherza AL, Axen K, Fain R, Hoffman RS, Dunbar CC, Haas F. Effect of naloxone on perceived exertion and exercise capacity during maximal cycle ergometry. *J Appl Physiol*. 2002;93(6):2023–8.
48. Robertson CV, Marino FE. A role for the prefrontal cortex in exercise tolerance and termination. *J Appl Physiol*. 2016;120(4):464–6.
49. Saanijoki T, Nummenmaa L, Koivumäki M, Löyttyniemi E, Kalliokoski KK, Hannukainen JC. Affective adaptation to repeated SIT and MICT protocols in insulin-resistant subjects. *Med Sci Sports Exerc*. 2018;50(1):18–27.
50. Saanijoki T, Nummenmaa L, Eskelinen JJ, et al. Affective responses to repeated sessions of high-intensity interval training. *Med Sci Sports Exerc*. 2015;47(12):2604–11.
51. Petruzzello SJ, Hall EE, Ekkekakis P. Regional brain activation as a biological marker of affective responsiveness to acute exercise: influence of fitness. *Psychophysiology*. 2001;38(1):99–106.
52. Hallgren MÅ, Moss ND, Gastin P. Regular exercise participation mediates the affective response to acute bouts of vigorous exercise. *J Sports Sci Med*. 2010;9(4):629–37.
53. Hoffman MD, Hoffman DR. Exercisers achieve greater acute exercise-induced mood enhancement than nonexercisers. *Arch Phys Med Rehabil*. 2008;89(2):358–63.
54. Chen YC, Chen C, Martínez RM, Etner JL, Cheng Y. Habitual physical activity mediates the acute exercise-induced modulation of anxiety-related amygdala functional connectivity. *Sci Rep*. 2019;9(1):19787.
55. Rhodes RE, Kates A. Can the affective response to exercise predict future motives and physical activity behavior? A systematic review of published evidence. *Ann Behav Med*. 2015;49(5):715–31.
56. Jackson AS, Sui X, Hébert JR, Church TS, Blair SN. Role of lifestyle and aging on the longitudinal change in cardiorespiratory fitness. *Arch Intern Med*. 2009;169(19):1781–7.
57. Lin X, Zhang X, Guo J, et al. Effects of exercise training on cardiorespiratory fitness and biomarkers of cardiometabolic health: a systematic review and meta-analysis of randomized controlled trials. *J Am Heart Assoc*. 2015;4(7):e002014.
58. Hoffmann P, Terenius L, Thorén P. Cerebrospinal fluid immunoreactive beta-endorphin concentration is increased by voluntary exercise in the spontaneously hypertensive rat. *Regul Pept*. 1990;28(2):233–9.
59. Su CF, Chang YY, Pai HH, Liu IM, Lo CY, Cheng JT. Mediation of beta-endorphin in exercise-induced improvement in insulin resistance in obese Zucker rats. *Diabetes Metab Res Rev*. 2005;21(2):175–82.
60. Shyu BC, Andersson SA, Thorén P. Endorphin mediated increase in pain threshold induced by long-lasting exercise in rats. *Life Sci*. 1982;30(10):833–40.
61. Brito RG, Rasmussen LA, Sluka KA. Regular physical activity prevents development of chronic muscle pain through modulation of supraspinal opioid and serotonergic mechanisms. *Pain Rep*. 2017;2(5):e618.
62. Bement MKH, Sluka KA. Low-intensity exercise reverses chronic muscle pain in the rat in a naloxone-dependent manner. *Arch Phys Med Rehabil*. 2005;86(9):1736–40.

63. Bruehl S, Burns JW, Koltyn K, et al. Are endogenous opioid mechanisms involved in the effects of aerobic exercise training on chronic low back pain? A randomized controlled trial. *Pain*. 2020;161(12):2887–97.
64. Bruehl S, Burns JW, Koltyn K, et al. Does aerobic exercise training alter responses to opioid analgesics in individuals with chronic low back pain? A randomized controlled trial. *Pain*. 2021;162(8):2204–13.
65. Raichlen DA, Foster AD, Seillier A, Giuffrida A, Gerdeman GL. Exercise-induced endocannabinoid signaling is modulated by intensity. *Eur J Appl Physiol*. 2013;113(4):869–75.
66. Koltyn KF, Brellenthin AG, Cook DB, Sehgal N, Hillard C. Mechanisms of exercise-induced hypoalgesia. *J Pain*. 2014;15(12):1294–304.
67. Brellenthin AG, Crombie KM, Hillard CJ, Koltyn KF. Endocannabinoid and mood responses to exercise in adults with varying activity levels. *Med Sci Sports Exerc*. 2017;49(8):1688–96.
68. Marin Bosch B, Bringard A, Logrieco MG, et al. A single session of moderate intensity exercise influences memory, endocannabinoids and brain derived neurotrophic factor levels in men. *Sci Rep*. 2021;11:14371.
69. Belitardo de Oliveira A, de Mello MT, Tufik S, Peres MFP. Weight loss and improved mood after aerobic exercise training are linked to lower plasma anandamide in healthy people. *Physiol Behav*. 2019;201:191–7.
70. Oliveira AB, Ribeiro RT, Mello MT, Tufik S, Peres MFP. Anandamide is related to clinical and cardiorespiratory benefits of aerobic exercise training in migraine patients: a randomized controlled clinical trial. *Cannabis Cannabinoid Res*. 2019;4(4):275–84.
71. Schoffelmeer AN, Hogenboom F, Wardeh G, De Vries TJ. Interactions between CB1 cannabinoid and mu opioid receptors mediating inhibition of neurotransmitter release in rat nucleus accumbens core. *Neuropharmacology*. 2006;51(4):773–81.
72. Wenzel JM, Cheer JF. Endocannabinoid regulation of reward and reinforcement through interaction with dopamine and endogenous opioid signaling. *Neuropsychopharmacology*. 2018;43(1):103–15.
73. Kantonen T, Karjalainen T, Pekkarinen L, et al. Cerebral μ -opioid and CB1 receptor systems have distinct roles in human feeding behavior. *Transl Psychiatry*. 2021;11:442.
74. Nummenmaa L, Tuominen L, Dunbar R, et al. Social touch modulates endogenous μ -opioid system activity in humans. *Neuroimage*. 2016;138:242–7.
75. Gunn RN, Lammertsma AA, Hume SP, Cunningham VJ. Parametric imaging of ligand-receptor binding in PET using a simplified reference region model. *Neuroimage*. 1997;6(4):279–87.
76. Manninen S, Karjalainen T, Tuominen LJ, et al. Cerebral grey matter density is associated with neuroreceptor and neurotransmitter availability: a combined PET and MRI study. *Neuroimage*. 2021;235:117968.
77. Heijnen S, Hommel B, Kibele A, Colzato LS. Neuromodulation of aerobic exercise—a review. *Front Psychol*. 2015;6:1890.

AUTHOR QUERIES

AUTHOR PLEASE ANSWER ALL QUERIES

AQ1 = Please check if authors name are correctly captured for given names (in red) and surnames (in blue) for indexing after publication.

AQ2 = Please provide the highest academic degree(s) of the corresponding author, if available.

AQ3 = Please expand VBM (mentioned only once).

AQ4 = Please expand MNI.

AQ5 = Please check if FDR has been defined correctly.

END OF AUTHOR QUERIES