

Physiology of the BOLD Signal


TURKU PET CENTRE NEUROIMAGING COURSE 2024


3.9.2024 Kerttu Seppälä

PhD Student, Turku PET Centre

kerttu.seppala@utu.fi

What is fMRI?





8:30-9:15 Physiology of the BOLD signal and T2* image acquisition

9:15-10:00 Experimental designs for fMRI

10:00-10:45 Preprocessing with fMRIPrep

10:45-12:00 Lunch break

12:00-12:45 General Linear Model

12:45-13:30 First level models for fMRI

13:30-13:45 Coffee break


13:45-14:30 Second level models for fMRI

14:30-15:15 Region of interest analysis



15:15-16:00 Data visualization

What is MRI and fMRI?

- Images taken with MRI scanner (Magnetic Resonance Imaging)
- **Anatomical MRI:** anatomical image gives structure (but they cannot reveal short-term physiological changes)
- **fMRI:** functional images
 - measure changes in subjects' brains function while they are focusing on a task in a MRI scanner or try not to do anything
 - Purpose
 - 1) finding specific part of the brain where mental process happens
 - 2) patterns of brain activation associated with mental processes
 - signal based on rapid changes in blood oxygenation over time on specific areas
 - the signal is coming from nuclear level, fMRI measurements noninvasive and are not interfering with neuronal firing or blood flow



Raw fMRI data, 1 volume





Collect enough volumes over time




Preprocessing
+ statistics



Beautifull fMRI images



How to collect data?





Siemens MAGNETOM Sola 1.5 T

<https://www.siemens-healthineers.com>

Siemens MAGNETOM Vida 3 T



Philips Ingenia 1.5 T

<https://www.philips.fi/>

Philips Ingenia 3 T



GE SIGNA 1.5 T

www.gehealthcare.com/products/magnetic-resonance-imaging/

GE SIGNA 3 T






Prepare the subject on the table
Some sites ask the subjects to change clothes, some not, please follow the rules on your own laboratory

While scanning:

- Noise → double ear protection
- Claustrophobia → communicate
- SAR (specific absorption rate) → no patients with fever
- Careful screening: tattoos, implants, ear rings, etc.



Place the subject into the scanner


Important: make the subject feel comfortable



Double hearing protection must be



Extra super important: the subject cannot move!
There are selection of different pillows to gently prevent subjects' head from moving



Screen for showing stimuli

Mirror to see the stimuli

Panic button sitting on a easy-reach place

Hands holding the buttons for tasks

Head in head coil, ears protected

Nice and comfortable
positioning, the subject is
ready to start the task

The magnet is always on



Safety first! The magnet is always on

- 1.5 T is 30 000 times stronger than earth magnetic field

IMPORTANT INSTRUCTIONS

Before entering the MR environment or MR system room, you must remove all metallic objects including hearing aids, dentures, partial plates, keys, beeper, cell phone, eyeglasses, hair pins, barrettes, jewelry, body piercing jewelry, watch, safety pins, paperclips, money clip, credit cards, bank cards, magnetic strip cards, coins, pens, pocket knife, nail clipper, tools, clothing with metal fasteners, & clothing with metallic threads.

Please consult the MRI Technologist or Radiologist if you have any question or concern **BEFORE** you enter the MR system room.



<https://www.harveynorman.com.au/amable-office-chair-red.html>

**True for everyone
and everything
entering the
scanner room**



WARNING: Certain implants, devices, or objects may be hazardous to you and/or may interfere with the MR procedure (i.e., MRI, MR angiography, functional MRI, MR spectroscopy). Do not enter the MR system room or MR environment if you have any question or concern regarding an implant, device, or object. Consult the MRI Technologist or Radiologist **BEFORE** entering the MR system room. The MR system magnet is **ALWAYS** on.

Please indicate if you have any of the following:

- | | | |
|------------------------------|-----------------------------|--|
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | Aneurysm clip(s) |
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | Cardiac pacemaker |
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | Implanted cardioverter defibrillator (ICD) |
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | Electronic implant or device |
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | Magnetically-activated implant or device |
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | Neurostimulation system |
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | Spinal cord stimulator |
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | Internal electrodes or wires |
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | Bone growth/bone fusion stimulator |
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | Cochlear, otologic, or other ear implant |
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | Insulin or other infusion pump |
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | Implanted drug infusion device |
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | Any type of prosthesis (eye, penile, etc.) |
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | Heart valve prosthesis |
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | Eyelid spring or wire |
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | Artificial or prosthetic limb |
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | Metallic stent, filter, or coil |
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | Shunt (spinal or intraventricular) |
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | Vascular access port and/or catheter |
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | Radiation seeds or implants |
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | Swan-Ganz or thermodilution catheter |
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | Medication patch (Nicotine, Nitroglycerine) |
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | Any metallic fragment or foreign body |
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | Wire mesh implant |
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | Tissue expander (e.g., breast) |
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | Surgical staples, clips, or metallic sutures |
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | Joint replacement (hip, knee, etc.) |
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | Bone/joint pin, screw, nail, wire, plate, etc. |
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | IUD, diaphragm, or pessary |
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | Are you here for an MRI examination? |
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | Dentures or partial plates |
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | Tattoo or permanent makeup |
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | Body piercing jewelry |
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | Hearing aid |
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | Other implant _____ |
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | Breathing problem or motion disorder |

(Remove before entering MR system room)





Fig. 2 – (a) Linear redness and swelling on her bilateral thighs after magnetic resonance imaging. (b) After a week, the redness gradually became blistering eruptions.

I attest that the above information is correct to the best of my knowledge. I read and understand the contents of this form and had the opportunity to ask questions regarding the information on this form and regarding the MR procedure that I am about to undergo.


Signature of Person Completing Form: _____

Signature


Date ____/____/____



Artifacts in T2w anatomical images on thighs.




a) Linear redness and swelling on patient's thigh after MRI.
b) After a week, the redness become blisters



a) Reddened area corresponded to the vertical lined on jogging pants
b) no visible metallic material; light, thin metal fibers were used in the pants.

Tokue H, Tokue A, Tsushima Y. **Unexpected magnetic resonance imaging burn injuries from jogging pants.** Radiol Case Rep. 2019 Sep 5;14(11):1348-1351.




When you know what to do, image acquisition is fun!


Where is the MRI signal coming from?

MR imaging is easy

- 1) **Place the subject** into the MRI scanner in strong magnetic field
- 2) Sent a **radio wave in**
- 3) Turn the **radio wave off**
- 4) The **subject emits a signal** for head coil to catch
- 5) **Reconstruct** the image




[Huettel Scott A.](#), [Song Allen W.](#), [McCarthy Gregory](#): Functional Magnetic Resonance Imaging, 2014, [Oxford University Press Inc](#)




Sprawls, Magnetic Resonance Imaging,
Online Edition, provided by Sprawls
Educational Foundation
<http://www.sprawls.org>



1



2



3



1. Spin:

- Intrinsic property
- Proton spinning on their axis
- 1) because proton has positive charge, spin generates electrical current on its surface
 - Magnetic moment in external magnetic field (μ)
- 2) because proton has odd-numbered atomic mass and when spinning → moving mass results in angular momentum (J)

2. Precession

- Spins align within external magnetic field
- Orange spins on lower energy state = parallel state (the amount always more!)
- Blue spins on higher energy state = antiparallel state

3. Net Magnetization (M):

- Difference between the number of spins in the parallel state and in the antiparallel state
- The more spins in the parallel state, the larger the M



THE MAGNETIC FIELD

Magnetic
Resonance
Imaging


Principles, Methods, and Techniques



Perry Sprawls



Sprawls



A) A neutral situation:

Spins are in external magnetic field, orange lower energy state, blue higher energy state


B) A radio wave comes in:

Some of the lower energy state spins jumps to higher energy state (orange spin into blue)

C) Incoming radio wave stops, outgoing starts:

Some of the high-energy state spins return to the lower energy state and release the absorbed energy as a radiofrequency wave with the same frequency as the excitation pulse

In conceptual way of thinking: 90° excitation pulse tilts the Net magnetization







Figure 3.12 A conceptual overview of T_1 recovery. The net magnetization tips into the transverse plane as a result of the absorption of energy by some spins (i.e., those that have changed from low- to high-energy states). Once the excitation pulse ceases, spins begin to release energy back into the surrounding environment. This causes the net magnetization to recover along the longitudinal axis, often returning to near its original amplitude within a few seconds (red dashed line).





Figure 3.11 A conceptual overview of T_2 decay. After the net magnetization has been tipped into the transverse plane, it rapidly decays because of a loss of coherence among the spins. For most types of tissue, the net magnetization available to generate the MR signal decays to near zero within a few hundred milliseconds (red dashed line).

Contrast

Contrast in medical imaging:
The intensity difference
between different quantities
(tissues)



Case Western Reserve University
Magnetic Resonance Imaging (MRI) of the Brain and Spine: Basics
<https://case.edu/med/neurology/NR/MRI%20Basics.htm>

T1 relaxation and contrast







Figure 4

Farrall, Magnetic Resonance Imaging, Practical Neurology 2006; 6: 318-325



Case Western Reserve University
Magnetic Resonance Imaging (MRI) of the Brain and Spine: Basics
<https://case.edu/med/neurology/NR/MRI%20Basics.htm>

T2 relaxation and contrast









Figure 15



Farrall, Magnetic Resonance Imaging, Practical Neurology 2006; 6: 318-325



Case Western Reserve University
 Magnetic Resonance Imaging (MRI) of the Brain and Spine: Basics
<https://case.edu/med/neurology/NR/MRI%20Basics.htm>



Local magnetic field
inhomogeneities →
 T_2^* contrast (even
faster decay)



$$\mu = \gamma J$$

$$\frac{d\mu}{dt} = \gamma (\mu \times B_0)$$

$$M = \frac{\Delta E}{2k_B T} n \mu_z \mathbf{z}$$

$$\omega_{\text{rot}} = \gamma B_{\text{leff}} = \gamma B_1 \quad \text{emf} = -i\omega_0 \int_V \overline{B}_1 \cdot M(t) dv$$

$$\frac{d\mathbf{M}}{dt} = \gamma \mathbf{M} \times \mathbf{B} + \frac{1}{T_1} (M_0 - M_z) - \frac{1}{T_2} (M_x + M_y)$$

Quantitative path

Where is the fMRI signal coming from?

Brain is full of arteries, capillaries and veins

- **Arteries** (oxygen rich blood from heart) → **capillaries** (exchange of oxygen to carbon dioxide) → **veins** (back to lungs)
- 100 billion neuros, 20 billion within cortex
- 800 mL / min of blood through average 1400 g brain = 15%-20 % of the blood flow in human body. Brain takes 2-3% of the body weight but **requires 20% of blood oxygen**
- Oxygen travels in hemoglobin
- 4 oxygen molecules in each hemoglobin molecule, 280 million hemoglobin molecules in each red blood cell




FIG. 1. Drawing of the cortical pial vessels. Right hemisphere. Female, 50 years. RED: Tributaries of the middle cerebral artery. GREEN: Tributaries of the anterior cerebral artery. BLUE: Tributaries of the posterior cerebral artery. Veins are shown in black.

Duvernoy HM, Delon S, Vannson JL. Cortical blood vessels of the human brain. Brain Res Bull. 1981 Nov;7(5):519-79. doi: 10.1016/0361-9230(81)90007-1. PMID: 7317796.







FIG. 60. Temporal pole (SEM). Male, 66 years. General view of the cortical vessels. (1) Cortical arteries. (2) Cortical vein. (3) Medullary artery (type A6). (4) Subcortical white matter ($\times 40$).

Duvernoy HM, Delon S, Vannson JL. Cortical blood vessels of the human brain. *Brain Res Bull.* 1981 Nov;7(5):519-79. doi: 10.1016/0361-9230(81)90007-1. PMID: 7317796.

fMRI and BOLD


- fMRI does not measure neuronal activity!
- fMRI measures physiological changes correlated with neuronal activity
- BOLD: **B**lood-**O**xygenation-**L**evel **D**ependent contrast
 - Activity of neurons increase metabolic requirements; blood becomes briefly deoxygenated
 - Oxygen travels in hemoglobin
 - Vascular system provides glucose and oxygen refill: arterial supply of oxygenated hemoglobin increases; the amount of the oxygenated blood increases on broad area more than needed
 - amount of deoxygenated hemoglobin decreases compared to the normal conditions
 - measure with T_2^*
 - brighter MRI



↑neural activity → ↑ blood flow → ↑ oxyhemoglobin → ↑ T_2^* → ↑ MR signal

Effect of Hemoglobin in Magnetic Field

OxyHb: diamagnetic
→ weak field inhomogeneities




Slower dephasing
slower T_2^*

Marijke Fagan-Endres,
"Fundamental studies
of heap leaching
hydrology using
magnetic resonance
imaging, Theses, Jan
2013,
DOI:10.17863/CAM.66
92


Uniform magnetic field






Paramagnetic material




Diamagnetic material



Ferromagnetic material





DeoxyHb: paramagnetic
→ strong field inhomogeneities



Fast dephasing
Fast T_2^*

BOLD signal and Hemodynamic
Response?

HDR: Hemodynamic Response



The change in the MRI signal triggered by neuronal activity

The **initial dip**

- Might reflect the decreased oxygenation before arteries provide more
- Is not detected in every study; Easier to detect on higher field (≥ 7 T)
- Allen Elster: Questions and Answers in MRI Website: BOLD and Brain Activity, <http://mriquestions.com/does-boldbrain-activity.html>:
"mechanism remains disputed: a) increased early metabolic extraction of blood oxygen, and/or b) increased local cerebral blood volume."

The positive **dominant peak**

- Maximal amplitude of HDR
- The overcompensation of used oxygen from arteries to neurons
- Slow signal: 4-6 s delay

The **post-stimulus undershoot**

- Happens most likely because combination of reduced blood flow and increased blood volume
- Easier to detect in block of multiple consecutive events





Fig. 7.10

The Actual Measured Signal

Changes in BOLD activation after presenting single event stimuli for subject from a voxel



Example of BOLD hemodynamic response to a hand squeezing task






Fig. 7.12

The same data as on the left but the timeseries has cut and organized timewise

Spatial Resolution

- Spatial Resolution: Ability to distinguish different locations within an image


Figure 1.7 The human brain at different spatial resolutions. Spatial resolution refers to the ability to resolve small differences in an image. In general, we can define spatial resolution based on the size of the elements (i.e., voxels) used to construct the image. The images shown here present the same brain sampled at five different element sizes: 8 mm (A); 4 mm (B); 2 mm (C); 1.5 mm (D); and 1 mm (E). Note that the gray-white structure is well represented in the latter three images, all of which were produced using element sizes that were less than half the typical gray matter thickness of 5 mm.





Spatial Resolution 2

- Structural images voxels maybe 1 x 1 x 1 mm
- Functional images voxels maybe 3 x 3 x 3 mm (depends on the question)
- BOLD signal is direct measure of the amount of deoxyhemoglobin in a voxel
- Partial volume effects: combination of different tissue types within a voxel (effect from large arteries / small capillaries)
- → Spatial smoothing for statistics and better signal-to-noise ratio

MRI



fMRI




Voxel: 3D volume element

30 slices, 64 x 64 voxels per slice → 122800 voxels

Table 7.1 Spatial Scales in the Human Brain


Structure	Scale (mm)
Brain	100
Gyri	10
Dominance column	1



Structure	Scale (mm)
Neuron	0.01
Synapse	0.001
Ion channel	0.00001

Temporal Resolution

- Determined by TR and by limitations of vascular system
 - **TR** = time of repetition (time for a volume)
 - HDR rises and falls within 10-15 s
 - **Duration** of the stimulus does not necessarily correspond with duration of neuronal activity
- fMRI is slow
 - neuronal activity is short $< 1s$
 - no snapshot of neuronal activity but an estimate of slower changes in vascular system
- Good TR?
 - Depending on the experiment (0,5 s – 3 s)
 - Smaller TR
 - more accurate estimation of HDR shape;
not necessary effect on amplitude








Figure 7.20 Effects of sampling rate (TR) on the measured hemodynamic response. In each figure, an idealized hemodynamic response is sampled at a different rate.

Timeseries



Every voxel



Many volumes over time




Summary

- fMRI:
 - neuroimaging technique with MRI scanners
 - measures physiological changes correlated with neuronal activity
 - Safe when used correctly, no ionizing radiation
- BOLD: Blood-Oxygenation-Level Dependent contrast

↑neural activity → ↑ blood flow → ↑ oxyhemoglobin → ↑ T2* → ↑ MR signal

- HDR: Hemodynamic Response



Question Time!